

Report on the observed climate, projected climate, and projected biodiversity changes for *Bergsåsen* under differing levels of warming

Jeff Price<sup>1</sup>, Nicole Forstenhäusler<sup>1</sup>, Erin Graham<sup>2</sup>,  
Timothy J. Osborn<sup>3</sup>, and Rachel Warren<sup>1</sup>

Report of the Wallace Initiative, July 1, 2024

<sup>1</sup>Tyndall Centre for Climate Change Research,  
School of Environmental Sciences, University of East Anglia, Norwich, UK.

<sup>2</sup>eResearch Centre, James Cook University, Townsville, Australia.

<sup>3</sup>Climatic Research Unit, School of Environmental Sciences, University of East Anglia, Norwich, UK.

## Contents

<b>Acknowledgements</b>	<b>3</b>
<b>Executive Summary</b>	<b>4</b>
Supporting Information . . . . .	7
<b>Overview</b>	<b>7</b>
<b>Climate</b>	<b>7</b>
Average Monthly High Temperature (usually the temperature of mid- to late-afternoon) .	8
Average Monthly Temperature . . . . .	9
Average Monthly Low Temperature (usually the temperature just before dawn) . . . . .	10
Precipitation . . . . .	11
Drought/Waterlogging . . . . .	12
<b>Population</b>	<b>13</b>
<b>Landcover changes</b>	<b>13</b>
<b>Biodiversity</b>	<b>14</b>
Local Extinction Risk . . . . .	14
Species Richness Remaining . . . . .	15
Refugia . . . . .	25
Adaptation Effort . . . . .	35
<b>More Detailed Information</b>	<b>44</b>
Climate . . . . .	44
Climate Variables . . . . .	45
Population Data . . . . .	47
Landcover Changes . . . . .	48
Biodiversity . . . . .	48
Developing robust adaptation plans in the light of climate projection uncertainties . . . . .	50
<b>Literature Cited</b>	<b>51</b>

## Acknowledgements

This work draws on nearly a decade of collaborative research between the Tyndall Centre for Climate Change Research, University of East Anglia, United Kingdom, and the eResearch Centre at James Cook University, Australia. We wish to thank those who played a major role in producing the original underlying models, data, and webpages used in the Wallace Initiative - J. VanDerWal, I. Atkinson, D. Baird, and P. Lawrey. We would also like to acknowledge the High Performance Computing Team at the University of East Anglia. Portions of this work have been funded from the MacArthur Foundation funding to World Wildlife Fund-US and IUCN (Ecosystems and Livelihoods Adaptation Network), WWF-UK (Wildlife in a Warming World), UK Natural Environment Research Council (NERC) grant no. NE/P014992/1 (IMPALA).

The climate projections were produced using the ClimGen system funded by NERC (QUEST-GSI; NE/E001831/1) and EU (HELIX; 603864) and we thank Ian Harris and Craig Wallace for producing the underlying data. We acknowledge the climate modelling groups, the Program for Climate Model Diagnosis and Intercomparison and the World Climate Research Programme's Working Group on Coupled Modelling for producing and making available the CMIP5 model output.

Parts of this work was funded with support from the European Union's Horizon 2020 research and innovation programmes under grant agreement No 101003687 (PROVIDE).










Figure ES1: Percent overall biodiversity remaining at 1 km resolution.



## Average Monthly High Temperature (usually the temperature of mid- to late-afternoon)

Table 1: Observed Average Monthly High Temperature (°C) with a comparison of the amount of change occurring between 1961-1990 and 1991-2020. Warmest refers to the warmest year in the 30-year period, coolest to the coolest year. In the warmest column, yellow shading indicates a temperature equal to that occurring approximately one in every three years ( $> 1$  SD) compared to 1961-1990; red shading indicates a temperature equal to or greater than that occurring one in every twenty years ( $>2$  SD) compared to 1961-1990. One way of looking at this is these years could be seen as “proxies” for what an average year in the future looks like.

Month	1961-1990			1991-2020			Difference 91-20 to 61-90
	Coolest	Average	Warmest	Coolest	Average	Warmest	Average
Jan	-9.2	-3.5	3.5	-6.0	-1.5	2.1	2.0
Feb	-10.5	-2.6	3.4	-6.9	-1.6	1.4	1.0
Mar	-3.2	0.3	3.7	-2.3	1.2	3.9	0.9
Apr	1.2	3.8	5.7	1.8	5.2	8.2	1.4
May	7.2	9.9	13.0	7.4	10.1	13.9	0.2
Jun	11.8	14.7	18.6	11.1	14.6	19.2	-0.1
Jul	13.9	15.7	18.5	14.3	16.8	19.6	1.1
Aug	12.5	14.9	19.9	13.2	15.9	19.6	1.0
Sep	8.2	10.4	12.9	9.2	11.7	13.7	1.3
Oct	2.6	5.9	9.5	1.4	6.0	8.6	0.1
Nov	-3.5	0.2	3.7	-3.9	1.3	5.5	1.0
Dec	-9.3	-1.9	3.0	-6.4	-0.6	3.4	1.3

Table 2: Projected Changes in Average Monthly High Temperature (°C) - Yellow shading indicates when the new average is equal to that occurring approximately one in every three years ( $> 1$  SD) compared to 1961-1990; red shading indicates when the new average is equal to or greater than that occurring one in every twenty years ( $>2$  SD) compared to 1961-1990.

Month	Scenario					
	1.5 °C	2 °C	2.5 °C	3 °C	3.5 °C	4 °C
Jan	1.5	2.2	2.9	3.6	4.2	4.9
Feb	1.4	2.0	2.6	3.2	3.9	4.5
Mar	1.3	1.9	2.5	3.1	3.7	4.3
Apr	1.7	2.4	3.1	3.8	4.6	5.3
May	1.6	2.3	3.0	3.7	4.4	5.1
Jun	1.3	1.8	2.4	3.0	3.5	4.1
Jul	1.3	1.8	2.3	2.9	3.4	4.0
Aug	1.4	2.0	2.6	3.2	3.8	4.4
Sep	1.4	2.0	2.6	3.2	3.8	4.4
Oct	1.5	2.1	2.7	3.4	4.0	4.6
Nov	1.6	2.3	3.1	3.8	4.5	5.2
Dec	1.6	2.3	3.0	3.7	4.4	5.1



## Average Monthly Temperature

Table 3: Observed Average Monthly Temperature (°C) with a comparison of the amount of change occurring between 1961-1990 and 1991-2020. Warmest refers to the warmest year in the 30-year period, coolest to the coolest year. In the warmest column, yellow shading indicates a temperature equal to that occurring approximately one in every three years ( $> 1$  SD) compared to 1961-1990; red shading indicates a temperature equal to or greater than that occurring one in every twenty years ( $> 2$  SD) compared to 1961-1990. One way of looking at this is these years could be seen as “proxies” for what an average year in the future looks like.

Month	1961-1990			1991-2020			Difference 91-20 to 61-90
	Coolest	Average	Warmest	Coolest	Average	Warmest	Average
Jan	-12.6	-6.4	1.0	-8.8	-4.1	-0.4	2.3
Feb	-13.6	-5.7	1.6	-10.6	-4.5	-0.3	1.2
Mar	-8.2	-3.1	0.8	-6.6	-2.1	0.9	1.0
Apr	-2.1	0.5	2.4	-1.4	1.7	4.1	1.2
May	3.8	6.1	8.7	4.2	6.3	9.3	0.2
Jun	7.9	10.4	13.1	7.4	10.5	14.3	0.1
Jul	10.0	11.8	14.1	10.7	12.9	15.5	1.1
Aug	9.2	11.2	15.0	10.2	12.2	14.9	1.0
Sep	4.7	7.4	9.6	6.0	8.6	10.6	1.2
Oct	0.2	3.6	7.6	-1.1	3.7	6.7	0.1
Nov	-6.1	-2.0	1.9	-6.3	-0.8	3.3	1.2
Dec	-12.1	-4.5	1.0	-9.5	-2.9	1.3	1.6

Table 4: Projected Changes in Average Monthly Temperature (°C) - Yellow shading indicates when the new average is equal to that occurring approximately one in every three years ( $> 1$  SD) compared to 1961-1990; red shading indicates when the new average is equal to or greater than that occurring one in every twenty years ( $> 2$  SD) compared to 1961-1990.

Month	Scenario					
	1.5 °C	2 °C	2.5 °C	3 °C	3.5 °C	4 °C
Jan	1.9	2.7	3.5	4.4	5.2	6.0
Feb	1.7	2.5	3.2	4.0	4.8	5.5
Mar	1.6	2.3	3.0	3.7	4.4	5.1
Apr	1.6	2.3	3.0	3.7	4.4	5.1
May	1.6	2.2	2.9	3.6	4.3	5.0
Jun	1.3	1.9	2.5	3.1	3.6	4.2
Jul	1.3	1.8	2.4	2.9	3.5	4.0
Aug	1.4	2.0	2.6	3.2	3.8	4.4
Sep	1.4	2.0	2.6	3.2	3.8	4.4
Oct	1.5	2.2	2.8	3.5	4.1	4.8
Nov	1.9	2.7	3.5	4.3	5.1	5.9
Dec	1.9	2.8	3.6	4.4	5.2	6.1

### Average Monthly Low Temperature (usually the temperature just before dawn)

Table 5: Observed Average Monthly Low Temperature (°C) with a comparison of the amount of change occurring between 1961-1990 and 1991-2020. Warmest refers to the warmest year in the 30-year period, coolest to the coolest year. In the warmest column, yellow shading indicates a temperature equal to that occurring approximately one in every three years ( $> 1$  SD) compared to 1961-1990; red shading indicates a temperature equal to or greater than that occurring one in every twenty years ( $>2$  SD) compared to 1961-1990. One way of looking at this is these years could be seen as “proxies” for what an average year in the future looks like.

Month	1961-1990			1991-2020			Difference 91-20 to 61-90
	Coolest	Average	Warmest	Coolest	Average	Warmest	Average
Jan	-16.1	-9.4	-1.2	-11.7	-6.8	-2.9	2.6
Feb	-16.8	-8.8	-0.2	-14.4	-7.4	-2.0	1.4
Mar	-13.3	-6.5	-2.1	-10.9	-5.4	-2.1	1.1
Apr	-5.9	-2.8	-0.8	-4.8	-1.7	0.2	1.1
May	0.4	2.3	4.5	0.8	2.6	4.9	0.2
Jun	3.6	6.2	8.7	3.8	6.4	9.4	0.2
Jul	6.1	8.0	10.4	7.1	9.2	11.7	1.2
Aug	6.0	7.5	10.2	6.5	8.6	10.9	1.1
Sep	0.9	4.5	6.8	2.8	5.5	7.6	1.0
Oct	-2.2	1.3	5.7	-3.6	1.3	4.8	0.0
Nov	-8.8	-4.3	0.2	-8.7	-2.8	1.4	1.5
Dec	-14.9	-7.2	-1.0	-12.7	-5.2	-0.8	2.0

Table 6: Projected Changes in Average Monthly Low Temperature (°C) - Yellow shading indicates when the new average is equal to that occurring approximately one in every three years ( $> 1$  SD) compared to 1961-1990; red shading indicates when the new average is equal to or greater than that occurring one in every twenty years ( $>2$  SD) compared to 1961-1990.

Month	Scenario					
	1.5 °C	2 °C	2.5 °C	3 °C	3.5 °C	4 °C
Jan	2.2	3.2	4.2	5.1	6.1	7.1
Feb	2.1	3.0	3.9	4.8	5.7	6.6
Mar	2.0	2.9	3.8	4.7	5.5	6.4
Apr	1.9	2.7	3.5	4.3	5.1	5.9
May	1.6	2.3	3.0	3.7	4.4	5.1
Jun	1.4	2.0	2.6	3.2	3.8	4.4
Jul	1.3	1.9	2.5	3.1	3.6	4.2
Aug	1.4	2.1	2.7	3.3	3.9	4.6
Sep	1.4	2.1	2.7	3.3	3.9	4.5
Oct	1.7	2.4	3.1	3.8	4.5	5.2
Nov	2.2	3.1	4.1	5.0	6.0	6.9
Dec	2.3	3.3	4.3	5.2	6.2	7.2

## Precipitation

Table 7: Observed Average Monthly Precipitation (mm/month) with a comparison of the amount of change occurring between 1961-1990 and 1991-2020. Wettest refers to the wettest year in the 30-year period, driest to the driest year. In the wettest and driest columns, yellow shading indicates precipitation amounts that exceed 1 SD above (wettest year) or below (driest year) the 1961-1990 average, i.e. amounts comparable to those that occur approximately one in every three years during 1961-1990; red shading indicates precipitation amounts occurring one in every twenty years compared to 1961-1990 ( $> 2$  SD for wettest year,  $< 2$  SD for driest). One way of looking at this is these years could be seen as “proxies” for what an average year in the future looks like.

Month	1961-1990			1991-2020			Difference 91-20 to 61-90
	Wettest	Average	Driest	Wettest	Average	Driest	Average
Jan	249.9	108.2	12.7	216.9	114.8	8.6	6.6
Feb	150.8	86.9	29.4	150.0	92.0	16.2	5.1
Mar	215.8	89.4	22.3	151.9	95.8	46.4	6.5
Apr	109.5	68.0	33.5	114.3	66.8	22.2	-1.2
May	97.8	53.8	25.9	86.5	58.4	28.3	4.6
Jun	167.3	77.6	23.6	141.3	88.7	36.8	11.1
Jul	152.8	102.7	26.3	180.4	93.1	28.3	-9.6
Aug	168.6	94.5	24.9	185.4	109.0	64.4	14.6
Sep	226.0	136.3	36.2	207.2	119.8	35.9	-16.5
Oct	222.5	133.1	58.0	196.5	124.3	56.8	-8.9
Nov	176.0	102.5	46.5	173.3	101.5	24.8	-1.1
Dec	320.1	127.9	49.8	264.7	130.9	34.6	3.0

Table 8: Projected Average Monthly Precipitation Change (mm) - Yellow shading indicates when the new average is equal to that occurring approximately one in every three years ( $> 1$  SD) compared to 1961-1990; red shading indicates when the new average is equal to or greater than that occurring one in every twenty years ( $> 2$  SD) compared to 1961-1990.

Month	Scenario					
	1.5 °C	2 °C	2.5 °C	3 °C	3.5 °C	4 °C
Jan	7.1	10.1	13.2	16.3	19.3	22.4
Feb	3.0	4.3	5.6	6.9	8.2	9.5
Mar	4.4	6.3	8.2	10.1	12.0	14.0
Apr	4.4	6.3	8.2	10.1	12.0	13.9
May	5.1	7.4	9.6	11.8	14.1	16.3
Jun	3.4	4.8	6.3	7.7	9.2	10.6
Jul	5.3	7.6	9.9	12.2	14.5	16.8
Aug	5.6	8.0	10.5	12.9	15.3	17.7
Sep	9.2	13.2	17.1	21.1	25.1	29.1
Oct	11.2	16.1	21.0	25.8	30.7	35.6
Nov	7.7	11.1	14.5	17.8	21.2	24.6
Dec	7.5	10.8	14.1	17.3	20.6	23.9

## Drought/Waterlogging

The drought metric used here measures severe meteorological drought (SPEI12, -1.5). It is the metric often used when looking at potential drought issues for agricultural and natural lands. The metric looks at droughts developing over the preceding 12 months before the 'counting' begins. Thus, an area identified as having a maximum drought duration of 12 months has been in drought for up to 24 months. The values in the table are calculated for the 30-year period for the observed or warming level given. Waterlogging is the reverse of the drought metric (SPEI12, +1.5) and is an indication of areas having excess moisture for extended periods, potentially leading to waterlogged soils.

Table 9: Observed number of months in severe drought or waterlogged in a 30-year period with a comparison of the amount of change occurring between 1961-1990 and 1986-2015.

	1961-1990	1986-2015	Difference 86-15 to 61-90
In drought	13	6.8	-6.2
Waterlogged	35	5.8	-29.2

Table 10: Observed maximum number of consecutive months in severe drought or waterlogged in a 30-year period with a comparison of the amount of change occurring between 1961-1990 and 1986-2015.

	1961-1990	1986-2015	Difference 86-15 to 61-90
In drought	6	6	0
Waterlogged	10	6	-4

Table 11: Changes in number of months in severe drought or waterlogged in a 30-year period.

	1.5 °C	2 °C	2.5 °C	3 °C	3.5 °C	4 °C
In drought	-2.6	-3.3	-3.7	-3.8	-3.7	-2.7
Waterlogged	9.6	13.4	17.5	22.0	26.6	30.5

Table 12: Changes in maximum number of consecutive months in severe drought or waterlogged in a 30-year period.

	1.5 °C	2 °C	2.5 °C	3 °C	3.5 °C	4 °C
In drought	-0.5	-1.0	-1.4	-1.5	-1.7	-1.6
Waterlogged	1.0	1.7	2.5	3.1	4.2	4.8

## Population

Table 13: Projected population for the years 2010 through 2100 at a 1 km<sup>2</sup> spatial resolution. These data are provided both in terms of the population within the protected area boundary, and those within an area including a 15 km wide buffer zone around the boundary. The data from 2000 and 2010 are interpolations of observed population sizes, the other periods are projections of future change in a 'middle-of-the-road' scenario with historical patterns of development continued through the 21st century.

Area	2000	2010	2030	2050	2070	2090	2100
Within region	5	4	5	6	6	7	7
Region plus buffer	892	779	926	1,056	1,200	1,267	1,271

## Landcover changes

Table 14: Percent landcover in 1992 and 2020, and change in landcover (300 m resolution). These figures are provided to assist in understanding how landcover has changed over time as this may have had immediate biodiversity implications in the area.

Landcover class	% in 1992	% in 2020	change (%)
Cropland, rainfed	1.89	1.89	0.00
Herbaceous cover	1.89	1.89	0.00
Mosaic cropland (>50%)/natural vegetation (tree, shrub, herbaceous cover) (< 50%)	1.89	3.77	1.88
Mosaic natural vegetation (tree, shrub, herbaceous cover) (>50%)/cropland (< 50%)	9.43	9.43	0.00
Tree cover, broadleaved, deciduous, closed to open (>15%)	3.77	3.77	0.00
Tree cover, needleleaved, evergreen, closed to open (>15%)	50.94	75.47	24.53
Tree cover, mixed leaf type (broadleaved and needleleaved)	0.00	3.77	3.77
Shrub or herbaceous cover, flooded, fresh/saline/brackish water	30.19	0.00	-30.19

## Biodiversity

The biodiversity information presented comes from models projecting climate suitability for ~135 000 terrestrial fungi, plants, invertebrates, and vertebrates. Resolution ~1 km<sup>2</sup>.

### Local Extinction Risk

Table 15: Percentage of species in different taxonomic groups projected to be at risk of local extinction owing to changes in climate alone. Yellow shading indicates areas projected to become climatically unsuitable for >25% of the species studied (by group); orange shading indicates areas projected to become climatically unsuitable for >50% of the species studied; and red shading indicates areas projected to become climatically unsuitable for >75% of the species studied. NA means there is insufficient data for that group in that area.

Taxa	1.5 °C	2 °C	3 °C	4 °C
<b>Biodiversity</b>	6.6	9.3	18.3	30.9
<b>Plants</b>	7.0	9.7	17.3	27.1
Ferns	6.2	7.9	12.7	19.3
Mosses	6.1	8.6	17.6	31.6
Pines	5.2	7.0	13.2	19.9
Flowering plants	7.5	10.3	18.1	28.6
Magnoliopsida	7.2	9.9	17.6	28.1
Liliopsida	8.2	11.0	18.4	28.7
Grasses	9.7	13.4	22.0	32.3
Lilies	8.2	10.6	14.5	22.7
Orchids	3.5	5.0	10.9	20.9
Palms	NA	NA	NA	NA
Vines	NA	NA	NA	NA
<b>Timber species</b>	1.5	2.5	5.2	10.8
<b>Animals</b>	8.4	11.8	22.7	37.9
Arthropoda	8.6	12.2	23.7	39.5
Arachnida	7.8	11.2	20.5	33.6
Spiders	8.0	11.3	20.6	34.1
<b>Insecta</b>	8.7	12.3	24.2	40.5
Bees	6.7	9.0	18.2	32.5
Beetles	10.1	14.2	28.5	47.8
True Bugs	9.7	13.3	26.2	44.1
Flies	10.8	14.6	24.0	37.6
Lepidoptera	5.0	6.9	17.7	36.0
Butterflies	4.9	7.0	14.0	25.7
Moths	5.0	7.0	18.0	36.6
Dragonflies	1.6	2.2	7.9	21.7
<b>Pollinators</b>	7.0	10.5	21.8	37.2
Chordata	3.5	4.2	9.0	16.1
<b>Amphibia</b>	2.8	6.2	14.9	28.6
<b>Aves</b>	2.9	3.4	7.4	13.5
<b>Mammals</b>	6.6	8.1	14.9	26.7
<b>Reptiles</b>	0.0	0.0	6.4	19.5

## Species Richness Remaining

Figures 1 to 9 show the average percent of the species (species richness) *remaining* within the boundaries of the area (also depicted on the map as a solid black line) for selected groups. This shows the spatial variability in the potential patterns of loss.





## Plants




Figure 2: Percent plants remaining at 1 km resolution.

## Amphibians




Figure 3: Percent amphibians remaining at 1 km resolution.

## Birds




Figure 4: Percent birds remaining at 1 km resolution.

## Mammals

## Reptiles




Figure 6: Percent reptiles remaining at 1 km resolution.

## Insects




Figure 7: Percent insects remaining at 1 km resolution.

## Pollinators




Figure 8: Percent pollinators remaining at 1 km resolution.

## Timber species




Figure 9: Percent timber species remaining at 1 km resolution.



## Refugia

Table 16 shows the percent of the area remaining a climatic refugium for different groups of species. Climatic refugia are defined as areas remaining climatically suitable for >75% of the species in each group. The two columns, for each warming level, are >0 (meaning at least one climate change model projects that the area is a refugium) and >10 (meaning that at least half of the models project an area is a refugium). The shading is – darker green, >75% of the area is a refugium; medium green, 50%-75% of the area is a refugium; light green, 25%-50% of the area is a refugium; and white, less than 25% of the area is a refugium.

Figures 10 to 17 show the number of climate models agreeing that a particular pixel (cell) is a refugium for the taxa indicated. These maps provide a spatial representation of the agreement in the models (or areas with potentially lower uncertainty) to be refugia for the different groups as well as how this potentially varies within the area under study.

Taxa	1.5 °C		2.0 °C		3.0 °C		4.0 °C	
	> 0	> 10	> 0	> 10	> 0	> 10	> 0	> 10
Biodiversity	100	100	100	100	100	99.9	100	68.6
Plants	100	100	100	100	100	100.0	100	74.0
Ferns	100	100	100	100	100	100.0	100	100.0
Mosses	100	100	100	100	100	100.0	100	74.0
Pines	100	100	100	100	100	100.0	100	100.0
Flowering plants	100	100	100	100	100	100.0	100	74.0
Magnoliopsida	100	100	100	100	100	100.0	100	74.0
Liliopsida	100	100	100	100	100	100.0	100	74.0
Grasses	100	100	100	100	100	100.0	100	68.6
Lilies	100	100	100	100	100	100.0	100	96.8
Orchids	100	100	100	100	100	100.0	100	100.0
Palms	NA	NA	NA	NA	NA	NA	NA	NA
Vines	NA	NA	NA	NA	NA	NA	NA	NA
Timber species	100	100	100	100	100	100.0	100	100.0
Animals	100	100	100	100	100	99.9	100	68.6
Arthropoda	100	100	100	100	100	99.9	100	68.6
Arachnida	100	100	100	100	100	100.0	100	68.6
Spiders	100	100	100	100	100	100.0	100	68.6
Insecta	100	100	100	100	100	99.9	100	68.6
Bees	100	100	100	100	100	100.0	100	74.0
Beetles	100	100	100	100	100	85.6	100	0.0
True Bugs	100	100	100	100	100	96.8	100	24.6
Flies	100	100	100	100	100	84.5	100	0.0
Lepidoptera	100	100	100	100	100	100.0	100	68.6
Butterflies	100	100	100	100	100	100.0	100	85.6
Moths	100	100	100	100	100	100.0	100	68.6
Dragonflies	100	100	100	100	100	100.0	100	100.0
Pollinators	100	100	100	100	100	100.0	100	68.6
Chordata	100	100	100	100	100	100.0	100	100.0
Amphibia	100	100	100	100	100	96.8	100	24.6
Aves	100	100	100	100	100	100.0	100	100.0
Mammals	100	100	100	100	100	100.0	100	84.5
Reptiles	100	100	100	100	100	100.0	100	100.0




Figure 10: Number of models in agreement for overall biodiversity refugia at 1 km resolution.

## Plants




Figure 11: Number of models in agreement for plant refugia at 1 km resolution.

## Amphibians




Figure 12: Number of models in agreement for amphibian refugia at 1 km resolution.

## Birds




Figure 13: Number of models in agreement for bird refugia.

## Mammals





Figure 14: Number of models in agreement for mammal refugia at 1 km resolution.





## Insects




Figure 16: Number of models in agreement for insect refugia at 1 km resolution.

## Pollinators




Figure 17: Number of models in agreement for pollinator refugia at 1 km resolution.






Figure 18: Adaptation effort for overall biodiversity at 1 km resolution.

## Plants




Figure 19: Adaptation effort for plants at 1 km resolution.

## Amphibians




Figure 20: Adaptation effort for amphibians at 1 km resolution.

## Birds




Figure 21: Adaptation effort for birds at 1 km resolution.

## Mammals




Figure 22: Adaptation effort for mammals at 1 km resolution.



## Reptiles




Figure 23: Adaptation effort for reptiles at 1 km resolution.

## Insects




Figure 24: Adaptation effort for insects at 1 km resolution.

## Pollinators




Figure 25: Adaptation effort for pollinators at 1 km resolution.

## More Detailed Information

### Climate

The following text reviews the observed climate, climate variability and potential climate changes in the region listed at the top of the tables. Text on the projected changes to biodiversity then follows. The climate output contains an analysis of the output of the Community Integrated Assessment System (CIAS; Warren et al., 2008) with downscaled climate change projections from the IPCC CMIP5 climate model patterns from the ClimGEN pattern scaling system (Osborn et al., 2016). Also provided are analyses of observed climate data from the Climatic Research Unit (CRU) TS 4.07 database (Harris et al., 2020).

This report looks at high, low, and average temperatures, precipitation, and meteorological drought/waterlogging (Price et al., 2022) for the area listed. These projected changes are explored in the context of the current climate variability to which the area is already exposed (1-2 standard deviations, shaded as yellow or red respectively). This assumes that many human and ecological systems may be largely resilient to changes laying within the bound of recently experienced natural variability, depending on the return rate of the event. The first set of tables (1 to 12) summarise observed climate variability and projected climate change. For most variables, comparisons are also provided between two recent time periods to show the current trends in climate in the area. The tables also provide a comparison between the magnitudes of projected climate changes with observed climate variability in terms of standard deviations (yellow - >1 standard deviation, occurring ~1 in 3 years; red is >2 standard deviations, occurring ~1 in 20 years). Thus, if a month is shaded as red it means that the future average climate is projected to exceed that currently occurring only once in every twenty years.

Both observational and projected climate change data presented here have a spatial resolution of 0.5° of latitude by 0.5° of longitude, all calculated monthly. The observed climate data comes from the University of East Anglia Climatic Research Unit CRU TS 4.07 dataset (Harris et al., 2020), which provides monthly gridded climate data through 2022. Versions of these data have previously been used extensively in IPCC reports, and in many different works on climate change impacts. The data presented here cover two time periods: 1961–1990 and 1991–2020 (except drought which uses 1986–2015). Summary statistics are provided giving the difference between the climates in 1961–1990 and 1991–2020 to provide information on what changes (if any) in temperature and precipitation have been observed between these two time periods. The first period, 1961–1990, is one of the standards used for climate modelling results and is a commonly used baseline for impact models (including the biodiversity results presented here); by this time, the world had warmed by 0.35°C since 1861–1890 (see HadCRUT4 dataset of Morice et al., 2012). By the second period (1990–2020) warming had increased to ~0.9–1.0°C since 1861–1890. Similarly, the IPCC (2023) states that global land temperatures were 1.59°C (1.34–1.83°C) warmer between 1850–1900 and 2011–2020.

### Future Climates

Projected climate data comes from the Community Integrated Assessment System (CIAS; Warren et al., 2008) and its component module ClimGen (Osborn et al., 2016). In this approach, a simple climate model is first used to project global temperature rises (using a probabilistic approach to encompass the key uncertainties in state-of-the-art global climate change projections) over the 21st century, as a time series. The Climatic Research Unit has a database of stored outputs from 21 general circulation models (GCM) from a model inter-comparison project known as CMIP5. These outputs provide the pattern of how climate variables are projected to change regionally for specific levels of global temperature rise. ClimGEN scales these patterns to the amount of warming provided





- ## Role of elevation and topography

## Drought

## Waterlogging

## Population Data

Price, J., Forstenhäusler, N., Graham, E., Osborn, T.J., and Warren, R. (2024) Report on the observed climate, projected climate, and projected biodiversity changes for *Bergsässen* under differing levels of warming. Report of the Wallace Initiative.

Shared Socioeconomic Pattern (SSP) 2 scenario. This is one of five scenarios used in the IPCC to look at alternative pathways for climate change. In the SSP2 scenario, the trends into the future basically follow those existing today; sometimes referred to as a 'middle of the road scenario'. A good general source on the different SSPs can be found at <https://www.carbonbrief.org/explainer-how-shared-socioeconomic-pathways-explore-future-climate-change/> with the underpinning science data discussed in Riahi et al. (2017). Different SSPs would therefore provide different projections of future populations. The projections themselves use a weighting where areas currently increasing in population continue to increase, and those that are decreasing continue to decrease. This is a major assumption that may or may not hold up over time. More information on the process used to derive the population projections and subsequent downscaling can be found in Jones and O'Neill (2016), and Gao (2017).

## Landcover Changes

Table 14 shows the percentage of different landcover types in 1992 and 2020 as well as the change between these two time periods within the boundaries of the area in this report. These data come from the European Space Agency Climate Change Initiative (Copernicus Climate Change Service, Climate Data Store, 2019; ESA, 2017) and have a resolution of 300 metres. The main limitation of the ESA data is that it is classified without supervision, and not 'ground-truthed'. This means that areas designated as 'forest' may be plantations of non-native species. These figures are provided to assist in understanding how these land cover classes have changed over time as this may have immediate biodiversity implications in the area studied for this report.

## Biodiversity

The biodiversity information presented here is from the Wallace Initiative. The Wallace Initiative modelled ~135 000 terrestrial fungi, plants, invertebrates, and vertebrates, at warming levels ranging from 1.5 °C to 6 °C, across 21 CMIP5 climate model patterns at a spatial resolution of ~20 km × 20 km based on occurrence data obtained from the Global Biodiversity Information Facility (GBIF.org, 2015). More information on the overall project, results, modelling methodology, caveats, and uses can be found in a series of papers (Jenkins et al., 2021; Price et al., 2024a; Saunders et al., 2023; Smith et al., 2018; Warren et al., 2018a,b, 2013). The data were also used for a number of figures and tables in Working Group II of the IPCC Sixth Assessment Report (AR6). The data used in these reports were then subsequently elevationally downscaled to ~1 km × 1 km (Saunders et al., 2023) to better understand which areas of each modelled 20 km cell or pixel might be lost sooner or persist longer. In short, a given 50 km or 20 km cell is an average of the temperatures for all elevations within that cell (i.e., the average elevation). In areas with a varied terrain, some areas will be warmer than the average and some will be cooler. Species in areas that are warmer than the average would be expected to potentially be more susceptible (exposed) to warming, while those in cooler areas would be expected to potentially be less susceptible (or be able to shift into these areas if they are currently too cool). Therefore, species within cooler areas within a climate 'cell' or 'pixel' would be expected to potentially be able to persist in that area longer.

## Local Extinctions (extirpations)

Table 15 shows the percentage of species in different taxa projected to be at risk of local extinction (extirpation, losses within the area of the report) owing to changes in climate alone. Yellow shaded areas are projected to become climatically unsuitable for >25% of the species studied (by taxa listed); orange areas are projected to become climatically unsuitable for >50% of the species studied; and red areas are projected to become climatically unsuitable for >75% of the species studied. NA means there is insufficient data in the cell to assess overall likelihoods. The climate suitability is the average change (ensemble of biodiversity models) across the 21 climate models examined.



## Species Richness Remaining

Figures 1 to 9 show the species richness remaining in each 1 km cell within the boundaries of the area under study (also depicted on the map as a solid black line) for selected taxa. This shows the spatial variability in the potential patterns of loss.

## Refugia

Table 16 shows the percent of the area remaining a climate refugia for the taxa. We define a climate refugia as an area remaining climatically suitable for >75% of the species in those taxa. The two columns, for each level of warming, are >0 (meaning at least one climate change model projects that the area is a refugia) and >10 (meaning that more than 10 models, out of 21, project the area remains a refugia). The shading is – darker green, >75% of the area is a refugia; medium green, 50–75% of the area is a refugia; light green, 25–50% of the area is a refugia; and white, less than 25% of the area is a refugia.

Figures 10 to 17 show the number of models in agreement that a particular pixel (cell) is a refugia for the taxa indicated. These maps provide a spatial representation of the agreement in the models (or areas with potentially lower uncertainty) to be refugia for the different taxa as well as how this potentially varies within the area under study.

The biodiversity refugia map is the minimum models in agreement between the plant and animal refugia.

## Adaptation Effort

Figures 18 to 25 present a spatial representation of the potential ‘adaptation effort’ that might be needed to maintain at least 75% of the species modelled (i.e., area remains climatically suitable) in each ~1km pixel. Adaptation effort is a combination of the number of climate models (+ 1 to 21) projecting an area is a refugia (remaining climatically suitable for >75% of the species) as well as the number of climate models (- 1 to -21) projecting the area to be an Area of Concern (becomes climatically unsuitable for >75% of the species) in each pixel. One way of looking at this is to consider areas with high values (+18 to +21) as being less exposed to climate change and thus potentially more resilient. Business-as-usual conservation, especially if coupled with building resilience around extreme climates (e.g., drought, heat waves) might be a reasonable adaptation approach to take. As the score drops, increasingly greater amounts of adaptation might be needed to maintain the existing species in that pixel. While micro-refugia (areas <1km) might be available, the amount of habitat available as micro-refugia would be less than the pixel. Once the adaptation effort drops into the negative zone, adaptation to maintain the existing species is likely to become increasingly difficult. At score of -15 to -21 the best approach might be to consider facilitating change as opposed to putting large efforts into trying to maintain existing species. Scores this low indicate that the area becomes climatically unsuitable for a large percentage of species. While this does not preclude micro-refugia, large areas (and potentially the area of conservation interest) would appear to be transforming. In the case of an area where restoration or reforestation is planned, then consideration might be given to planting the species that might be expected to move into the area, given enough time (considering species with similar structure and native, if possible). This type of adaptation begins to make the new ‘habitat’ that species from surrounding areas will need to autonomously adapt to climate change.

There are many complexities in these analyses. Not least of which is that an area may remain a refugia for vertebrates and yet potentially become unsuitable for many of the species making up the habitat or food resources for these species. If the habitat becomes unsuitable, or food becomes more unavailable then this is likely to have major implications for those taxa that a cell remains

a refugium for. With increasing warming, fewer areas remain refugia, more areas become areas of concern, and adaptation effort increases (i.e., becomes more negative).

### **Developing robust adaptation plans in the light of climate projection uncertainties**

Climate change adaptation experts recommend an iterative risk management approach, particularly where climate change projections or future vulnerability is uncertain. Conceptual approaches for prioritising potential adaptation options might include: (i) implementing low cost ‘no regret’ adaptation plans, such as removal of concomitant stresses; (ii) in areas where it is unclear whether drying or wetting is projected, creating adaptation plans relating to changes in management to incorporate future projected climate change that remain flexible (e.g., either to adaptively manage or plan for both wetting and drying, or to be able to switch rapidly from managing/planning for wetting to what is needed for drying). Since climate change generally includes increases in climate variability, even in a future wetter climate, there may still be more droughts. This implies that adaptation to changes in precipitation needs to incorporate flexibility on both long and short timescales to cater for both wetting and drying in areas where the sign of precipitation projection differs across models. Even in areas where the sign of precipitation change is consistent between models (e.g.~positive), increases in climate variability on shorter timescales may still imply a need to cater for increased short-term drying. (iii) avoiding implementing plans that lock in the system to being able to cater for only the present day climate, (thus ignoring warming) or catering only for wetting (when actually drying may occur).

## Literature Cited

- Copernicus Climate Change Service, Climate Data Store (2019). *Land Cover Classification Gridded Maps from 1992 to Present Derived from Satellite Observation*. Copernicus Climate Change Service (C3S) Climate Data Store (CDS). DOI: [10.24381/cds.006f2c9a](https://doi.org/10.24381/cds.006f2c9a).
- ESA (2017). *Land Cover CCI Product User Guide Version 2*. Technical Report. URL: [http://maps.elie.ucl.ac.be/CCI/viewer/download/ESACCI-LC-Ph2-PUGv2\\_2.0.pdf](http://maps.elie.ucl.ac.be/CCI/viewer/download/ESACCI-LC-Ph2-PUGv2_2.0.pdf).
- Gao, J. (2017). *Downscaling Global Spatial Population Projections from 1/8-Degree to 1-Km Grid Cells*. NCAR Technical Note NCAR/TN-537+STR. DOI: [10.5065/D60Z721H](https://doi.org/10.5065/D60Z721H).
- GBIF.org (2015). *GBIF Occurrence Download*. Version March 2015. The Global Biodiversity Information Facility. DOI: [10.15468/DL.KECDHX](https://doi.org/10.15468/DL.KECDHX). URL: <https://www.gbif.org/occurrence/download/0000129-150523225239109>.
- Harris, I., Osborn, T. J., Jones, P., and Lister, D. (2020). Version 4 of the CRU TS Monthly High-Resolution Gridded Multivariate Climate Dataset. *Scientific Data* 7 (1) (1), pp. 1–18. DOI: [10/ggq4fw](https://doi.org/10/ggq4fw).
- IPCC (2013). *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Ed. by T. F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P. M. Midgley. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press. 1535 pp.
- IPCC (2023). Summary for Policymakers. In: *Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Ed. by Core Writing Team, H. Lee, and J. Romero. Geneva, Switzerland: IPCC, pp. 1–34. DOI: [10.59327/IPCC/AR6-9789291691647.001](https://doi.org/10.59327/IPCC/AR6-9789291691647.001).
- Jenkins, R. L. M., Warren, R. F., and Price, J. T. (2021). Addressing Risks to Biodiversity Arising from a Changing Climate: The Need for Ecosystem Restoration in the Tana River Basin, Kenya. *PLOS ONE* 16 (7), e0254879. DOI: [10.1371/journal.pone.0254879](https://doi.org/10.1371/journal.pone.0254879).
- Jones, B. and O'Neill, B. C. (2016). Spatially Explicit Global Population Scenarios Consistent with the Shared Socioeconomic Pathways. *Environmental Research Letters* 11 (8), p. 084003. DOI: [10.1088/1748-9326/11/8/084003](https://doi.org/10.1088/1748-9326/11/8/084003).
- Morice, C. P., Kennedy, J. J., Rayner, N. A., and Jones, P. D. (2012). Quantifying Uncertainties in Global and Regional Temperature Change Using an Ensemble of Observational Estimates: The HadCRUT4 Data Set. *Journal of Geophysical Research: Atmospheres* 117 (D8). DOI: [10.1029/2011JD017187](https://doi.org/10.1029/2011JD017187).
- Osborn, T. J., Wallace, C. J., Harris, I. C., and Melvin, T. M. (2016). Pattern Scaling Using ClimGen: Monthly-Resolution Future Climate Scenarios Including Changes in the Variability of Precipitation. *Climatic Change* 134 (3), pp. 353–369. DOI: [10.1007/s10584-015-1509-9](https://doi.org/10.1007/s10584-015-1509-9).
- Price, J., Warren, R., and Forstnhäusler, N. (2024a). Biodiversity Losses Associated with Global Warming of 1.5 to 4 °C above Pre-Industrial Levels in Six Countries. *Climatic Change* 177 (3), p. 47. DOI: [10.1007/s10584-023-03666-2](https://doi.org/10.1007/s10584-023-03666-2).
- Price, J., Warren, R., Forstnhäusler, N., Jenkins, R., and Graham, E. (2024b). Assessing the Potential Risks of Climate Change on the Natural Capital of Six Countries Resulting from Global Warming of 1.5 to 4 °C above Pre-Industrial Levels. *Climatic Change* 177 (3), p. 46. DOI: [10.1007/s10584-023-03650-w](https://doi.org/10.1007/s10584-023-03650-w).
- Price, J., Warren, R., Forstnhäusler, N., Wallace, C., Jenkins, R., Osborn, T. J., and Van Vuuren, D. P. (2022). Quantification of Meteorological Drought Risks between 1.5 °C and 4 °C of Global Warming in Six Countries. *Climatic Change* 174 (1), p. 12. DOI: [10.1007/s10584-022-03359-2](https://doi.org/10.1007/s10584-022-03359-2).
- Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O'Neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaresma, J. C., KC, S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K., Hasegawa, T., Havlik, P., Humpenöder, F., Da Silva, L. A., Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaat, D., Masui, T., Rogelj,

- J., Strefler, J., Drouet, L., Krey, V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L., Doelman, J. C., Kainuma, M., Klimont, Z., Marangoni, G., Lotze-Campen, H., Obersteiner, M., Tabeau, A., and Tavoni, M. (2017). The Shared Socioeconomic Pathways and Their Energy, Land Use, and Greenhouse Gas Emissions Implications: An Overview. *Global Environmental Change* 42, pp. 153–168. DOI: [10.1016/j.gloenvcha.2016.05.009](https://doi.org/10.1016/j.gloenvcha.2016.05.009).
- Saunders, S., Grand, J., Bateman, B., Meek, M., Wilsey, C., Forstenhaeusler, N., Graham, E., Warren, R., and Price, J. (2023). Integrating Climate-Change Refugia into 30 by 30 Conservation Planning in North America. *Frontiers in Ecology & the Environment* 21 (2). DOI: [10.1002/fee.2592](https://doi.org/10.1002/fee.2592).
- Smith, P., Price, J., Molotoks, A., Warren, R., and Malhi, Y. (2018). Impacts on Terrestrial Biodiversity of Moving from a 2°C to a 1.5°C Target. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 376 (2119). DOI: [10.1098/rsta.2016.0456](https://doi.org/10.1098/rsta.2016.0456).
- UNEP-WCMC and IUCN (2024). *The World Database on Protected Areas (WDPA) and World Database on Other Effective Area-based Conservation Measures (WD-OECM)*. Version March 2024. Cambridge, UK: UNEP-WCMC and IUCN. URL: [www.protectedplanet.net](http://www.protectedplanet.net).
- Warren, R., Price, J., Graham, E., Forstenhaeusler, N., and VanDerWal, J. (2018a). The Projected Effect on Insects, Vertebrates, and Plants of Limiting Global Warming to 1.5°C Rather than 2°C. *Science* 360 (6390), pp. 791–795. DOI: [10.1126/science.aar3646](https://doi.org/10.1126/science.aar3646).
- Warren, R., de la Nava Santos, S., Arnell, N. W., Bane, M., Barker, T., Barton, C., Ford, R., Fussell, H. -M., Hankin, R. K. S., Klein, R., Linstead, C., Kohler, J., Mitchell, T. D., Osborn, T. J., Pan, H., Raper, S. C. B., Riley, G., Schellnhüber, H. J., Winne, S., and Anderson, D. (2008). Development and Illustrative Outputs of the Community Integrated Assessment System (CIAS), a Multi-Institutional Modular Integrated Assessment Approach for Modelling Climate Change. *Environmental Modelling & Software* 23 (5), pp. 592–610. DOI: [10.1016/j.envsoft.2007.09.002](https://doi.org/10.1016/j.envsoft.2007.09.002).
- Warren, R., Price, J., VanDerWal, J., Cornelius, S., and Sohl, H. (2018b). The Implications of the United Nations Paris Agreement on Climate Change for Globally Significant Biodiversity Areas. *Climatic Change* 147 (3), pp. 395–409. DOI: [10.1007/s10584-018-2158-6](https://doi.org/10.1007/s10584-018-2158-6).
- Warren, R., VanDerWal, J., Price, J., Welbergen, J. A., Atkinson, I., Ramirez-Villegas, J., Osborn, T. J., Jarvis, A., Shoo, L. P., Williams, S. E., and Lowe, J. (2013). Quantifying the Benefit of Early Climate Change Mitigation in Avoiding Biodiversity Loss. *Nature Climate Change* 3 (7), pp. 678–682. DOI: [10.1038/nclimate1887](https://doi.org/10.1038/nclimate1887).
- WWF (2018). *Wildlife in a Warming World: The Effects of Climate Change on Biodiversity in WWF's Priority Places*. URL: [https://www.wwf.org.uk/sites/default/files/2018-03/WWF\\_Wildlife\\_in\\_a\\_Warming\\_World.pdf](https://www.wwf.org.uk/sites/default/files/2018-03/WWF_Wildlife_in_a_Warming_World.pdf).